# FULVI PRO® 10%

## **GUARANTEED ANALYSIS**

Derived from: Oxidized Leonardite. Delivered by: MICRO CARBON TECHNOLOGY®+ \*The standard for ascertaining Fulvic Acid percentage has not been established.

### ALSO CONTAINS NON-PLANT FOOD INGREDIENT:

6.0% Organic Matter (derived from leonardite)

## **DIRECTIONS FOR USE**

• Shake well before use. • Designed to be applied by soil application. • FULVI PRO® 10% supplies fulvic acid to crops and should be applied when soil analysis identifies the need and proper rate of application for this product. • Best results will be obtained when application is concentrated in the active root zone or when applied directly to the soil followed by a shallow cultivation. • Can be applied in combination with compatible plant growth regulators, pesticides, or other liquid fertilizers. • If compatibility is in question, jar test a small quantity. • Use caution when adding to high analysis/multi-nutrient solutions, highly acidic or basic solutions, or unstable suspensions. • Do not apply this product in concentrations greater than 10%. • Do not apply HUMA GRO® nutrients during the hottest time of the day or to plants under drought or flood stress. • Application timing, intervals and rates may vary according to individual crop requirement, stage of development, available nutrient levels in the soil, and overall nutritional status of the crop. • Applications can be made as often as every 15 to 30 days as needed. • Soil moisture is required for maximum bioactivity. If soil is dry, moisture should be provided by irrigation. • Consult your local HUMA GRO® Representative or other agricultural specialist for crop-specific recommendations. • Do not store this product below 10°C (50°F) or above 30°C (90°F).

This product is intended as a supplement to a regular fertilization program and will not, by itself, provide all the nutrients normally required by agricultural crops.

#### STORAGE AND DISPOSAL:

Keep product in original container. Do not transfer into food or drink containers. Triple rinse container when empty for recycling. Always dispose of container in accordance with local, state and/or federal regulations.

# **METHOD OF APPLICATION**

SUGGESTED RATE PER HECTARE Field Crops / Tree or Vine Crops

Foliar band application at 50% coverage	up to 2.5 liters /
Foliar broadcast or sprinklers: solid, set, linear, pivot (100% speed)	up to 5 liters / up to 10 liters
Soil banded or injected, through drip tape or micro sprinklers	up to 10 liters / up to 20 liters
Soil broadcast spray incorporated.	

#### CONDITIONS OF SALE:

flood, or furrow irrigated

The information contained on this label is believed to be accurate and reliable. Buyer and user acknowledge and assume all liability resulting from the use of this material. Follow directions carefully. Timing, method of application, weather, crop conditions and other factors are beyond the control of the seller.

#### KEEP OUT OF REACH OF CHILDREN

A clear to slightly hazy, amber liquid having a slight characteristic odor. The vapors, mists and liquid may cause severe irritation or burns to all tissues contacted. Phosphoric Acid may generate flammable Hydrogen gas on contact with most metals. The NIOSH I.D.L.H. for Phosphoric Acid is: 1,000 mg/m3.



CLASSIFICATION: SKIN CORROSION - CATEGORY 1A SIGNAL WORD: DANGER

HAZARD STATEMENT: H314 Causes severe skin burns and eve damage. PRECAUTIONARY STATEMENT: P260 Do not breathe dusts/mist/vapors; P280 Wear protective gloves/protective clothing/eye protection/face protection; P264 Wash hands thoroughly after handling.

1.05 kg per liter @ 20° C. 9.18 lb per gallon @ 68° F. pH 1.5 - 2.5

10 liters • Net Wt. 10.50 kg

LOT# 12345678



\*A Registered Trademark of and Produced by **BIO HUMA NETICS, INC.** © 2005-2020 1331 W. Houston Ave., Gilbert, AZ 85233 • 480.961.1220

> www.HumaGro.com HGI-620-07302020

