

LUCKY 7®

GUARANTEED ANALYSIS 7-7-7

| | |
|--|--------|
| Total Nitrogen (N) | 7.00% |
| 1.5% Ammoniacal Nitrogen | |
| 3.0% Urea Nitrogen | |
| 2.5% Nitrate Nitrogen | |
| Available Phosphate (P ₂ O ₅) | 7.00% |
| Soluble Potash (K ₂ O) | 7.00% |
| Magnesium (Mg) | 0.50% |
| Sulfur (S) | 1.00% |
| 1.00% Combined Sulfur (S) | |
| Boron (B) | 0.02% |
| Cobalt (Co) | 0.01% |
| Iron (Fe) | 0.10% |
| 0.10% Chelated Iron (Fe) | |
| Manganese (Mn) | 0.05% |
| 0.05% Chelated Manganese (Mn) | |
| Molybdenum (Mo) | 0.006% |
| Zinc (Zn) | 0.05% |
| 0.05% Chelated Zinc (Zn) | |

Derived from: Urea, Ammonium Nitrate, Phosphoric Acid, Potassium Hydroxide, Magnesium Sulfate, Boric Acid, Cobalt Chloride, Iron HEDTA, Manganese EDTA, Sodium Molybdate, Zinc EDTA. (Chelating agents are Hydroxyethylethylenediaminetriacetate and Ethylenediaminetetraacetate.)

ALSO CONTAINS NON-PLANT FOOD INGREDIENT:

7.0% Organic Matter (derived from leonardite)

DIRECTIONS FOR USE

- Shake well before use. • Designed to be applied by both soil and/or foliar application. • LUCKY 7® supplies nitrogen, phosphorus, potassium, and other nutrients to crops and should be applied when tissue and soil analysis identifies the need and proper rate of application for this product. • Best results will be obtained when application is concentrated in the active root zone or on the leaf surface. • Can be applied in combination with compatible plant growth regulators, pesticides, or other liquid fertilizers. • If compatibility is in question, jar test a small quantity. • Apply foliar sprays with sufficient water to ensure uniform coverage without running off leaf surfaces. • Do not apply this product foliarly in concentrations greater than 5%. • For maximum results, apply in the early morning or in the evening when air temperatures are optimum for foliar application. • Do not apply HUMA GRO® nutrients during the hottest time of the day or to plants under drought or flood stress. • Application timing, intervals and rates may vary according to individual crop requirement, stage of development, available nutrient levels in the soil, and overall nutritional status of the crop. • Applications can be made as often as every 7 to 10 days as needed. • Consult your local HUMA GRO® Representative or other agricultural specialist for crop-specific recommendations. • Do not store this product below 50°F (10°C) or above 90°F (30°C).

METHOD OF APPLICATION

SUGGESTED RATE PER ACRE
Field Crops / Tree or Vine Crops

| | |
|---|-----------------------------------|
| Foliar band application at 50% coverage | up to 1 quart / --- |
| Foliar broadcast or sprinklers: solid, set, linear, pivot (100% speed) | up to 2 quarts / up to 1 gallon |
| Soil banded or injected, through drip tape or micro sprinklers | up to 1 gallon / up to 2 gallons |
| Soil broadcast spray incorporated, flood, or furrow irrigated | up to 2 gallons / up to 4 gallons |

STORAGE AND DISPOSAL:

Keep product in original container. Do not transfer into food or drink containers. Triple rinse when empty for recycling. Always dispose of container in accordance with local, state and/or federal regulations.

This product is intended as a supplement to a complete fertilization program and will not, by itself, provide all the nutrients normally required by agricultural crops.

CONDITIONS OF SALE:

The information contained on this label is believed to be accurate and reliable. Buyer and user acknowledge and assume all liability resulting from the use of this material. Follow directions carefully. Timing, method of application, weather, crop conditions and other factors are beyond the control of the seller.

WARNING: Application of this material in excess may result in forage crops containing levels of Molybdenum (Mo) which are toxic to ruminant animals.

KEEP OUT OF REACH OF CHILDREN

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>

A clear to slightly hazy, acidic, purple to brown liquid having no characteristic odor. The liquid and mists may be irritating to the eyes and skin. Inhalation of mists may be irritating to the entire respiratory tract. Ingestion of this product may cause gastrointestinal irritation, cardiovascular and central nervous system effects.

CLASSIFICATION: HAZARD CATEGORY 3

SIGNAL WORD: WARNING

HAZARD STATEMENT: H316 - WARNING - causes mild skin irritation

PRECAUTIONARY STATEMENT: P332+P313 If skin irritation occurs, get medical attention/advice.

10.80 lb per gallon @ 68° F.
1.20 kg per liter @ 20° C.
pH 3.5 - 4.5

2.5 gallons • Net Wt. 27.00 lb
9.46 liters • Net Wt. 11.35 kg
LOT# 12345678



*Registered Trademarks of and Produced by

BIO HUMA NETICS, INC. © 2005-2020

1331 W. Houston Ave., Gilbert, AZ 85233 • 480.961.1220

www.HumaGro.com

HG-777-07292020

