

HUMA PRO[®] 16

GUARANTEED ANALYSIS

Total Humic Acids (HA)..... 16.00%*

*Analysis using the Colorimetric Method

Derived from: Oxidized Leonardite

Other Methods:

Total Humic Acids (HA)..... 10.00%[†]

[†]Analysis using the CDFA Method

Total Humic Acids (HA)..... 10.00%[‡]

[‡]Analysis using the HPTA or ISO 19822 Method

DIRECTIONS FOR USE

• Shake well before use. • Designed to be applied to the soil. • HUMA PRO[®] 16 supplies humic acids to crops and should be applied when soil analysis identifies the need and proper rate of application for this product. • Best results will be obtained when application is concentrated in the active root zone or when applied directly to the soil followed by a shallow cultivation. • Can be applied in combination with compatible plant growth regulators, pesticides, or granular fertilizers. • If compatibility is in question, jar test a small quantity. • Do not apply this product in concentrations greater than 10%. • Use caution when adding to high analysis/multi-nutrient solutions, highly acidic or basic solutions, or unstable suspensions. • Do not apply HUMA GRO[®] nutrients during the hottest time of the day or to plants under drought or flood stress. • Application timing, intervals and rates may vary according to individual crop requirement, stage of development, available nutrient levels in the soil, and overall nutritional status of the crop. • Applications can be made as often as every 15 to 30 days as needed. • Soil moisture is required for maximum bioactivity. If soil is dry, moisture should be provided by irrigation. • Consult your local HUMA GRO[®] representative or other agricultural specialist for crop-specific recommendations. • Do not store this product below 10°C (50°F) or above 30°C (90°F).

This product is intended as a supplement to a regular fertilization program and will not, by itself, provide the nutrients normally required by agricultural crops.

STORAGE AND DISPOSAL:

Keep product in original container. Do not transfer into food or drink containers. Triple rinse container when empty for recycling. Always dispose of container in accordance with local, state and/or federal regulations.

METHOD OF APPLICATION

SUGGESTED RATE PER HECTARE
Field Crops / Tree or Vine Crops

Soil banded, injected, side dress, drip tape, or micro sprinklers

up to 10 liters / up to 20 liters

Sprinklers: solid set, drag lines, linear, or pivot @ 100% speed

up to 20 liters / up to 40 liters

Soil broadcast spray incorporated, flood or furrow irrigated

up to 40 liters / up to 80 liters

CONDITIONS OF SALE:

The information contained on this label is believed to be accurate and reliable. Buyer and user acknowledge and assume all liability resulting from the use of this material. Follow directions carefully. Timing, method of application, weather, crop conditions and other factors are beyond the control of the seller.

KEEP OUT OF REACH OF CHILDREN

A dark brown to black, alkaline liquid having no characteristic odor. The liquid and mists may be irritating to the eyes and skin. Inhalation of mists may be irritating to the entire respiratory tract. Ingestion of this product may cause gastrointestinal irritation.

CLASSIFICATION: HAZARD CATEGORY 3 - MILD SKIN IRRITATION

SIGNAL WORD: WARNING

HAZARD STATEMENT: H316 - WARNING - Causes mild skin irritation.

PRECAUTIONARY STATEMENT: P332+P313 If skin irritation occurs, get medical attention/advice.

1.10 kg per liter @ 20° C.

9.01 lb per gallon @ 68° F.

pH 11.0 - 12.0

10 liters • Net Wt. 11.00 kg

LOT# 12345678

HUMA GRO[®]

*Registered Trademarks of and Produced by

BIO HUMA NETICS, INC. © 2005-2020

1331 W. Houston Ave., Gilbert, AZ 85233 • 480.961.1220

www.HumaGro.com

HGI-616-10292020

