

HUMA BURST® STABLESOL®

GUARANTEED ANALYSIS 0-0-20

Soluble Potash (K₂O) **20.0%**
Derived from: Potassium Hydroxide and Potassium Sulfate

Humic Acids **70.00%**
Determined using BaCl₂ Method

Derived from: Oxidized Leonardite

DIRECTIONS FOR USE

- HUMA BURST® STABLESOL® may promote the uptake of micronutrients and should be applied when soil analysis identifies the need and proper rate of application for this product.
- HUMA BURST® STABLESOL® is soluble over a pH range of 2.0 to 12.0. This unique feature allows HUMA BURST® STABLESOL® to be mixed and applied with most commercially available fertilizers and micronutrients.
- If compatibility is in question, jar test a small quantity.
- Best results will be obtained when application is concentrated in the active root zone or when applied directly to the soil followed by a shallow cultivation.
- Can be applied in combination with compatible plant growth regulators, pesticides, or other fertilizer solutions.
- Apply HUMA BURST® STABLESOL® by broadcast spray whether through fertigation, drip or spray machine. HUMA BURST® STABLESOL® can be applied to all soil types for ornamentals, lawns, gardens and landscapes.
- Avoid spreading during high temperatures or in times of drought.
- Application timing, intervals and rates may vary according to individual crop requirement, stage of development, available nutrient levels in the soil, and overall nutritional status of the crop.
- Applications can be made as often as every 15 days as needed.
- Soil moisture is required for maximum bioactivity. If soil is dry, moisture should be provided by irrigation.
- Consult your local HUMA GRO® representative or other agricultural specialist for crop-specific recommendations.

This product is intended as a supplement to a regular fertilization program and will not, by itself, provide all the nutrients normally required by agricultural crops.

STORAGE AND DISPOSAL:

Keep product in original bag. Do not transfer into food or drink containers. Always dispose of bag in accordance with local, state and/or federal regulations.

CROP TYPE

APPLICATION RATE

Vegetable crops, strawberries	Up to 2 lb/ac (2.25 kg/ha), diluted with 25 gal (100 l) of water At planting: Banded, side dressed, or injected in furrow At flower set and again 2 weeks prior to harvest: with irrigation
Row crops	Up to 3 lb/ac (3.36 kg/ha), diluted with 25 gal (100 l) of water At planting: Banded, side dressed, or injected in furrow At flower set and again 2 weeks prior to harvest: with irrigation
Citrus, Olives, Deciduous Fruit & Nut Trees	Up to 4 lb/ac (4.5 kg/ha), diluted with 25 gal (100 l) of water At flower set and again 2 weeks prior to harvest: with irrigation

CONDITIONS OF SALE:

The information contained on this label is believed to be accurate and reliable. Buyer and user acknowledge and assume all liability resulting from the use of this material. Follow directions carefully. Timing, method of application, weather, crop conditions and other factors are beyond the control of the seller.

KEEP OUT OF REACH OF CHILDREN

A brown powder having a mild characteristic odor. Ingestion of this product may cause gastrointestinal irritation or pain.

CLASSIFICATION: HAZARD CATEGORY 3 - MILD SKIN IRRITATION

SIGNAL WORD: WARNING

HAZARD STATEMENT: H316 - WARNING - Causes mild skin irritation.

PRECAUTIONARY STATEMENT: P332+P313 If skin irritation occurs, get medical attention/advice.

Net Wt. 55 lb

Net Wt. 25 kg

HUMA GRO®

*Registered Trademarks of and Produced by

BIO HUMA NETICS, INC. © 2018-2022

1331 W. Houston Ave., Gilbert, AZ 85233 • 480.961.1220

www.HumaGro.com

HG-650-06232022

